## **Post Test**

## Self-Study Module: Addressing Maternal and Neonatal Mortality

- 1. True or False? Current data shows maternal mortality is decreasing in the United States
  - a. True
  - b. False
- 2. The most common time for a woman to die in a pregnancy related death is:
  - a. During the pregnancy
  - b. Within 42 days of the birth
  - c. 42 days 1 year of the birth
- 3. What is the most common cause of death when assessing the overall causes of death related to pregnancy (for all maternal deaths)
  - a. Hemorrhage
  - b. Cardiovascular/Coronary Condition
  - c. Both
  - d. Neither
- 4. Patient Safety Bundles are an important piece of improving overall patient safety by
  - a. Providing a step by step process for hospitals to begin implementation of evidence based practices.
  - b. Utilizing Readiness, Recognition & Prevention, Response and Reporting/Systems Learning to address specific issues in maternal mortality.
  - c. Both
- 5. True or False? The New Maternal Deaths Acts of 2018 signed into law in December 2018, establishes support and some funding for review of maternal deaths through maternal mortality review committees at state levels. Committees will also provide recommendations for prevention of future maternal deaths.
  - a. True
  - b. False
- 6. True or False? The infant mortality rate in South Carolina decreased to 6.5 deaths per 1000 births in 2017?
  - a. True
  - b. False
- 7. True or False? The leading cause of infant mortality is SIDS?
  - a. True
  - b. False
- 8. True or False? Heart defects are known as a structural birth defect?
  - a. True
  - b. False
- 9. True or False? Hearing loss is the most common functional or developmental birth defect in South Carolina?
  - a. True
  - b. False
- 10. True or False? The leading cause of neonatal death is prematurity?
  - a. True
  - b. False